Indications:

Saniderm® Bandages can be used on abrasions, lacerations, new tattoos, and as a protective cover to prevent skin damage from friction.

Directions for use:

1. Cut the Saniderm bandage to size, rounding the corners and allowing at least a 1-inch border around the application area.

2. Clean the area with a soap mixture and dry thoroughly, ENSURE ALL RESIDUES HAVE BEEN REMOVED FROM THE SKIN. Lotions and Ointments are not necessary as they can interfere with adhesion.

3. Peel off the paper backing to expose the adhesive side.

4. Gently apply the center of the bandage to the area, using light pressure to smooth the film from the center out. Do not stretch the bandage or pull it tightly across the skin.

5. Using the perforations, remove the plastic top layer from the clear Saniderm film.

6. The first bandage should stay on for 8-24 hours and may fill up with blood and plasma. A second piece of Saniderm can then be worn for 3-5 days. If the wound weeps into the second piece of film, remove at the next 24-hour period and apply a 3rd piece. The third piece would then stay on for 5 days.

7. Saniderm® should not be worn for more than 7 days consecutively.

To remove Saniderm®

Gently pull the bandage down over itself slowly. Avoid pulling upward or ripping it off quickly. For easiest removal, use warm water to help loosen the adhesive.

Warnings:

Use only as directed. Do not apply to infected wounds. Do not cover the nose or mouth as it will prevent breathing. Saniderm® should be kept away from babies and children to avoid suffocation. • For external use only. This product contains adhesives and may cause allergic reactions in people who have adhesive allergies. • If allergic reaction or if any irritation occurs, discontinue use immediately, and if severe, consult a physician. • Saniderm® does not contain any UV protection and should be kept out of direct sunlight.

Please visit Saniderm.com

and click on our Knowledge Base for more information, FAQs, detailed instructions, videos, and helpful tips.

Saniderm Medcial LLC (626) 386-3376
PO Box 660 support@saniderm.com
Lehi, Utah 84043-1156 https://saniderm.com

STERILE EO

LATEX

40°F (5°C) 90°F (30°C)

CE 0197

EC REP

Shanghai International Holding Crop. GmbH (Europe) Eiffestrasse 80, 20537 Hamburg, Germany